

Tasigna[®] (nilotinib) (Oral)

Document Number: IC-0124

Last Review Date: 09/05/2023

Date of Origin: 11/01/2012

Dates Reviewed: 12/2012, 11/2013, 08/2014, 07/2015, 07/2016, 08/2017, 01/2018, 04/2018, 07/2018, 07/2019, 07/2020, 07/2021, 11/2021, 07/2022, 07/2023, 09/2023

I. Length of Authorization ¹

Coverage will be provided for 6 months and may be renewed.

- Patients with Ph+ CML-CP who have achieved a sustained molecular response should be evaluated for discontinuation after taking nilotinib for a minimum of 3 years. §
- For patients with possible TKI-resistant CML, coverage will be provided for 3 months and may be renewed for 6 months if the treatment response milestone criteria for TKI-sensitive disease is achieved (*Refer to Section IV for specific response criteria*)

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- Tasigna 50 mg capsules: 4 capsules per day
- Tasigna 150 mg capsules: 4 capsules per day
- Tasigna 200 mg capsules: 4 capsules per day

B. Max Units (per dose and over time) [HPCS Unit]:

- 800 mg per day

III. Initial Approval Criteria ¹

Coverage is provided in the following conditions:

- Patient is at least 18 years of age (unless otherwise specified); **AND**
- Patient has a baseline QTc interval of ≤ 480 ms **AND** does not have a history of long QT-syndrome; **AND**

Universal Criteria ¹

- Patient does not have hypokalemia or hypomagnesemia; **AND**
- Patient will avoid concomitant use of all of the following:

- Coadministration with strong CYP3A inhibitors (e.g., itraconazole, ketoconazole, clarithromycin, etc.), or if therapy is unavoidable, the patient will be monitored closely for adverse reaction and/or dose modifications will be implemented; **AND**
- Coadministration with strong CYP3A inducers (e.g., rifampin, carbamazepine, St. John's Wort, etc.); **AND**
- Coadministration with proton pump inhibitors (PPIs), or if acid-reduction therapy is required, H₂-receptor antagonists or antacids may be used at staggered administration times; **AND**
- Coadministration with drugs that prolong QT/QTc interval (e.g., amiodarone, sotalol, levofloxacin, venlafaxine, quetiapine, sumatriptan, etc.); **AND**

Chronic Myelogenous Leukemia (CML) † ‡ Φ 1,2,4,9-14

- Patient has Philadelphia chromosome-positive (Ph+) or *BCR::ABL1* positive disease; **AND**
- Patient does not have any of the following *BCR::ABL1* mutations: T315I, Y253H, E255K/V, or F359V/C/I (****NOTE:** This does not apply to newly diagnosed chronic phase disease or continued therapy); **AND**
 - Patient is resistant, intolerant, or had an inadequate response to prior therapy consisting of a 3 month trial or longer with a tyrosine kinase inhibitor (e.g., imatinib, bosutinib, dasatinib, etc.) †; **AND**
 - Patient is at least 1 year of age; **AND**
 - Used as a single agent for chronic phase or accelerated phase disease; **OR**
 - Used as primary treatment †; **AND**
 - Used as a single agent for newly diagnosed chronic phase disease in patients at least 1 year of age †; **OR**
 - Used as a single agent for accelerated phase disease; **OR**
 - Used as a single agent for myeloid blast phase disease if not a candidate for induction chemotherapy; **OR**
 - Used in combination with steroids for lymphoid blast phase disease if not a candidate for induction chemotherapy; **OR**
 - Used in combination with induction chemotherapy for disease in lymphoid blast phase or myeloid blast phase; **OR**
 - Used as switch therapy ‡; **AND**
 - Patient received primary therapy with one of the following: imatinib, bosutinib, or dasatinib; **AND**
 - Patient has *BCR::ABL1* transcript levels:
 - >0.1% to 1% at 12 months (if treatment goal is treatment-free remission); **OR**
 - >1% to 10% at 12 months; **OR**
 - >10% at any response milestone; **OR**

- Used as continued therapy ‡; **AND**
 - Patient has *BCR::ABL1* transcript levels:
 - ≤10% at any response milestone; **OR**
 - >10% at 3 months; **OR**
- Used post-allogeneic hematopoietic stem cell transplant (HCT) ‡; **AND**
 - Used for at least one year in patients with prior complete cytogenetic response (CCyR) for accelerated or blast phase disease; **OR**
 - Used as follow-up therapy in patients with molecular relapse (*BCR::ABL1* transcript positive) following CCyR; **OR**
 - Used as follow-up therapy in patients with relapse or less than CCyR; **OR**
- Re-initiation of treatment ‡; **AND**
 - Patient lost molecular response (MMR or MR4.0) after discontinuation of therapy with nilotinib

Acute Lymphoblastic Leukemia (ALL) ‡^{2,3,5}

- Patient has Philadelphia chromosome-positive (Ph+) B-ALL; **AND**
- Patient does not have any of the following *BCR::ABL1* mutations: T315I, Y253H, E255K/V, F359V/C/I or G250E; **AND**
 - Used for relapsed or refractory disease; **AND**
 - Used as a single agent; **OR**
 - Used in combination with inotuzumab ozogamicin; **OR**
 - Used in combination with blinatumomab; **OR**
 - Used as maintenance therapy; **AND**
 - Used in combination with POMP regimen (vincristine and prednisone with or without methotrexate and mercaptopurine); **AND**
 - Used following consolidation therapy for patients with negative minimal residual disease (if not already included in a multi-part regimen); **OR**
 - Used as single agent therapy; **AND**
 - Used post-hematopoietic stem cell transplant; **OR**
 - Used in patients unfit for additional therapies; **OR**
 - Used as induction therapy; **AND**
 - Used as frontline therapy OR for relapsed/refractory disease (if not previously given); **AND**
 - Used in combination with a corticosteroid; **OR**
 - Used in combination with vincristine and dexamethasone; **OR**
 - Used as a component of a multiagent chemotherapy regimen (*Note: May be used in patients at least 15 years of age*); **OR**

- Used as consolidation therapy; **AND**
 - Used for relapsed/refractory disease (if not previously given); **AND**
 - Used as a component of a multiagent chemotherapy regimen (*Note: May be used in patients at least 15 years of age*); **OR**
 - Used as frontline therapy; **AND**
 - Used as a component of a multiagent chemotherapy regimen (*Note: May be used in patients at least 15 years of age*); **OR**
 - Used as single agent therapy in patients unfit for additional therapies; **OR**
 - Used in combination with blinatumomab; **AND**
 - Used for persistent/rising minimal residual disease after complete response to induction therapy; **OR**
 - Used for negative minimal residual disease after complete response to induction therapy if patient is not a candidate for multiagent therapy

Gastrointestinal Stromal Tumors (GIST) ‡^{2,7,17}

- Used as a single agent; **AND**
- Patient has gross residual (R2 resection), unresectable primary, recurrent, or metastatic disease **OR** tumor rupture; **AND**
- Disease has progressed on imatinib, sunitinib, regorafenib, and ripretinib

Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Fusion Genes ‡^{2,15,16}

- Patient has eosinophilia and ABL1 rearrangement; **AND**
 - Patient has chronic or blast phase myeloid or lymphoid neoplasms; **AND**
 - Used as a single agent; **OR**
 - Patient has blast phase lymphoid, myeloid, or mixed lineage neoplasms; **AND**
 - Used in combination with ALL- or AML-type induction chemotherapy followed by allogeneic HCT (if eligible)

Soft Tissue Sarcoma (STS) ‡^{2,6}

- Patient has pigmented villonodular synovitis/tenosynovial giant cell tumor (PVNS/TGCT); **AND**
- Used as a single agent

Cutaneous Melanoma ‡²

- Used as subsequent therapy as a single agent; **AND**
- Patient has metastatic or unresectable disease with activating mutations of KIT; **AND**
- Used for disease progression, intolerance, and/or projected risk of progression with BRAF-targeted therapy

IV. Renewal Criteria^{1-4,6,15,17}

Coverage can be renewed based upon the following criteria:

- Patient continues to meet the universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in Section III; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: electrolyte abnormalities (hypophosphatemia, hypokalemia, hyperkalemia, hypocalcemia, hyponatremia), myelosuppression (neutropenia, thrombocytopenia, anemia), QT prolongation, cardiac and arterial vascular occlusive events, pancreatitis and elevated serum lipase, hepatotoxicity (severe changes in liver function tests), tumor lysis syndrome, hemorrhage, fluid retention, growth retardation in pediatric patients, etc. **AND**

Acute Lymphoblastic Leukemia (ALL)

- Treatment response or stabilization of disease as indicated by CBC, bone marrow cytogenic analysis, QPCR, or FISH

Chronic Myelogenous Leukemia (CML)

- Re-initiation of treatment:
 - Patient lost molecular response (MMR or MR4.0) after discontinuation of therapy with nilotinib; **OR**
- Treatment response as indicated by one of the following:
 - Patient has TKI-sensitive disease as confirmed by one of the following *BCR::ABL1* (IS) transcript levels:
 - ≤ 10% at 3 months or 6 months; **OR**
 - > 0.1% to 1% at 12 months and beyond (if treatment goal is long-term survival); **OR**
 - ≤ 0.1% at 12 months and beyond (if treatment goal is treatment-free remission); **OR**
 - Patient has possible TKI resistant disease as confirmed by a > 50% reduction in *BCR::ABL1* (IS) transcript levels compared to pretreatment baseline **OR** minimally above the 10% cutoff; **AND**
 - Treatment can be continued for an additional 3 months; **AND**
 - Follow-up *BCR::ABL1* (IS) transcript levels will be drawn after the 3 month extended treatment period to determine if continued treatment is warranted (i.e., patient meets the treatment response milestone criteria for TKI-sensitive disease as noted above)

Note: cytogenetic assessment of response may be used if quantitative RT-PCR (QPCR) using International Scale (IS) for *BCR::ABL1* is not available

Gastrointestinal Stromal Tumors (GIST)

- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread

Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Fusion Genes

- Disease response as evidenced by at least one of the following:
 - Decrease in spleen size or improvements in other myelofibrosis symptoms (such as weakness, fatigue, cough, dyspnea, myalgias, angioedema, rash, fever, rhinitis, etc.)
 - Stabilization or improvement as evidenced by a complete response [CR] (i.e. morphologic, cytogenetic or molecular complete response CR), complete hematologic response, or a partial response by CBC, bone marrow cytogenetic analysis, QPCR, or FISH

Soft Tissue Sarcoma (STS)

- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread

Cutaneous Melanoma

- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread

§ Consider discontinuation of treatment in patients with Ph+ CML-CP	
<p><u>Newly diagnosed Ph+ CML-CP who have:</u></p> <ul style="list-style-type: none"> – been treated with Tasigna for at least 3 years – maintained a molecular response of at least MR4.0 (corresponding to $\text{BCR-ABL/ABL} \leq 0.01\% \text{ IS}$) for one year prior to discontinuation of therapy – achieved an MR4.5 for the last assessment taken immediately prior to discontinuation of therapy – been confirmed to express the typical BCR-ABL transcripts (e13a2/b2a2 or e14a2/b3a2) – no history of accelerated phase or blast crisis – no history of prior attempts of treatment-free remission discontinuation that resulted in relapse. 	<p><u>Ph+ CML-CP that are resistant or intolerant to treatment with imatinib who have achieved a sustained molecular response (MR4.5) on Tasigna who have:</u></p> <ul style="list-style-type: none"> – been treated with Tasigna for a minimum of 3 years – been treated with imatinib only prior to treatment with Tasigna – achieved a molecular response of MR4.5 (corresponding to $\text{BCR-ABL/ABL} \leq 0.0032\% \text{ IS}$) – sustained an MR4.5 for a minimum of one year immediately prior to discontinuation of therapy – been confirmed to express the typical BCR-ABL transcripts (e13a2/b2a2 or e14a2/b3a2) – no history of accelerated phase or blast crisis – no history of prior attempts of treatment-free remission discontinuation that resulted in relapse.

V. Dosage/Administration ^{1,5,7,8,18-21}

Indication	Dose
Chronic Myelogenous Leukemia (CML)	<i>Adults with newly diagnosed chronic phase CML</i> 300 orally twice daily

	<p><u><i>Adults with resistant or intolerant chronic or accelerated phase CML</i></u></p> <p>400mg orally twice daily</p> <p><u><i>Pediatrics</i></u></p> <p>230 mg/m² orally twice daily (rounded to the nearest 50mg dose to a maximum single dose of 400 mg and up to a maximum daily dose of 800 mg)</p>
Acute Lymphoblastic Leukemia (ALL)	400 mg orally twice daily
Gastrointestinal Stromal Tumors (GIST)	400 mg orally twice daily
Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Fusion Genes	Up to 400 mg orally twice daily
Soft Tissue Sarcoma (STS)	400 mg orally twice daily
Cutaneous Melanoma	400 mg orally twice daily

VI. Billing Code/Availability Information

HCPCS Code:

- J8999: Prescription drug, oral, chemotherapeutic, not otherwise specified

NDC(s):

- Tasigna 50 mg capsule: 00078-0951-xx
- Tasigna 150 mg capsule: 00078-0592-xx
- Tasigna 200 mg capsule: 00078-0526-xx

VII. References

1. Tasigna [package insert]. East Hanover, NJ; Novartis Pharm. Corp; September 2021. Accessed August 2023.
2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for nilotinib hydrochloride monohydrate. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc.” To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed August 2023.
3. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for Acute Lymphoblastic Leukemia. 1.2023. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc.” To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2023.

4. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for Chronic Myeloid Leukemia 1.2024. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc.” To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed August 2023.
5. Kim DY, Joo YD, Lim SN, et al. Nilotinib combined with multiagent chemotherapy for newly diagnosed Philadelphia-positive acute lymphoblastic leukemia. *Blood* 2015; 126:746-756.
6. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for Soft Tissue Sarcoma 2.2023. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc.” To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2023.
7. Montemurro M, Schöffski P, Reichardt P, et al. Nilotinib in the treatment of advanced gastrointestinal stromal tumours resistant to both imatinib and sunitinib. *Eur J Cancer*. 2009;45(13):2293-2297. doi:10.1016/j.ejca.2009.04.030.
8. Sawaki A, Nishida T, Doi T, et al. Phase 2 study of nilotinib as third-line therapy for patients with gastrointestinal stromal tumor. *Cancer*. 2011;117(20):4633-4641. doi:10.1002/cncr.26120.
9. Saglio G, Kim DW, Issaragrisil S, et al. Nilotinib versus imatinib for newly diagnosed chronic myeloid leukemia. *N Engl J Med*. 2010;362(24):2251-2259. doi:10.1056/NEJMoa0912614.
10. Kantarjian HM, Giles FJ, Bhatta KN, et al. Nilotinib is effective in patients with chronic myeloid leukemia in chronic phase after imatinib resistance or intolerance: 24-month follow-up results. *Blood*. 2011;117(4):1141-1145. doi:10.1182/blood-2010-03-277152.
11. le Coutre PD, Giles FJ, Hochhaus A, et al. Nilotinib in patients with Ph+ chronic myeloid leukemia in accelerated phase following imatinib resistance or intolerance: 24-month follow-up results. *Leukemia*. 2012;26(6):1189-1194. doi:10.1038/leu.2011.323.
12. Ross DM, Masszi T, Gómez Casares MT, et al. Durable treatment-free remission in patients with chronic myeloid leukemia in chronic phase following frontline nilotinib: 96-week update of the ENESTfreedom study. *J Cancer Res Clin Oncol*. 2018;144(5):945-954. doi:10.1007/s00432-018-2604-x.
13. Mahon FX, Boquimpani C, Kim DW, et al. Treatment-Free Remission After Second-Line Nilotinib Treatment in Patients With Chronic Myeloid Leukemia in Chronic Phase: Results From a Single-Group, Phase 2, Open-Label Study. *Ann Intern Med*. 2018;168(7):461-470. doi:10.7326/M17-1094.

14. Hijiya N, Maschan A, Rizzari C, et al. Phase 2 study of nilotinib in pediatric patients with Philadelphia chromosome-positive chronic myeloid leukemia. *Blood*. 2019;134(23):2036-2045. doi:10.1182/blood.2019000069.
15. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Fusion Genes. Version 1.2023. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2023.
16. Metzgeroth G, Erben P, Martin H, Mousset S, et al. Limited clinical activity of nilotinib and sorafenib in FIP1L1-PDGFRα positive chronic eosinophilic leukemia with imatinib-resistant T674I mutation. *Leukemia* volume 26, pages 162–164 (2012).
17. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Gastrointestinal Stromal Tumors. Version 1.2023. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2023.
18. Gelderblom H, Cropet C, Chevreau C, et al. Nilotinib in locally advanced pigmented villonodular synovitis: a multicentre, open-label, single-arm, phase 2 trial. *Lancet Oncol*. 2018 May;19(5):639-648. doi: 10.1016/S1470-2045(18)30143-8.
19. Guo J, Carvajai R, Dummer R, et al. Efficacy and safety of nilotinib in patients with *KIT*^T mutated metastatic or inoperable melanoma: final results from the global, single-arm, phase II TEAM trial. *Annals of Oncology*. Volume 28, Issue 6, June 2017, Pages 1380-1387. <https://doi.org/10.1093/annonc/mdx079>.
20. Lee S, Kim T, Kim Y, et al. Phase II Trial of Nilotinib in Patients With Metastatic Malignant Melanoma Harboring KIT Gene Aberration: A Multicenter Trial of Korean Cancer Study Group (UN10-06). *Oncologist*. 2015 Nov;20(11):1312-9. doi: 10.1634/theoncologist.2015-0161. Epub 2015 Sep 30.
21. Schwaab J, Naumann N, Luebke J, et al. Response to tyrosine kinase inhibitors in myeloid neoplasms associated with PCM1-JAK2, BCR-JAK2 and ETV6-ABL1 fusion genes. *Clinical Trial Am J Hematol*. 2020 Jul;95(7):824-833. doi: 10.1002/ajh.25825. Epub 2020 Apr 28.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C43.0	Malignant melanoma of lip
C43.111	Malignant melanoma of right upper eyelid, including canthu

C43.112	Malignant melanoma of right lower eyelid, including canthus
C43.121	Malignant melanoma of left upper eyelid, including canthus
C43.122	Malignant melanoma of left lower eyelid, including canthus
C43.20	Malignant melanoma of unspecified ear and external auricular canal
C43.21	Malignant melanoma of right ear and external auricular canal
C43.22	Malignant melanoma of left ear and external auricular canal
C43.30	Malignant melanoma of unspecified part of face
C43.31	Malignant melanoma of nose
C43.39	Malignant melanoma of other parts of face
C43.4	Malignant melanoma of scalp and neck
C43.51	Malignant melanoma of anal skin
C43.52	Malignant melanoma of skin of breast
C43.59	Malignant melanoma of other part of trunk
C43.60	Malignant melanoma of unspecified upper limb, including shoulder
C43.61	Malignant melanoma of right upper limb, including shoulder
C43.62	Malignant melanoma of left upper limb, including shoulder
C43.70	Malignant melanoma of unspecified lower limb, including hip
C43.71	Malignant melanoma of right lower limb, including hip
C43.72	Malignant melanoma of left lower limb, including hip
C43.8	Malignant melanoma of overlapping sites of skin
C43.9	Malignant melanoma of skin, unspecified
C49.A0	Gastrointestinal stromal tumor, unspecified site
C49.A1	Gastrointestinal stromal tumor of esophagus
C49.A2	Gastrointestinal stromal tumor of stomach
C49.A3	Gastrointestinal stromal tumor of small intestine
C49.A4	Gastrointestinal stromal tumor of large intestine
C49.A5	Gastrointestinal stromal tumor of rectum
C49.A9	Gastrointestinal stromal tumor of other sites
C49.4	Malignant neoplasm of connective and soft tissue of abdomen
C49.5	Malignant neoplasm of connective and soft tissue of pelvis
C49.8	Malignant neoplasm of overlapping sites of connective and soft tissue
C49.9	Malignant neoplasm of connective and soft tissue, unspecified
C49.10	Malignant neoplasm of connective and soft tissue of unspecified upper limb, including shoulder
C49.11	Malignant neoplasm of connective and soft tissue of right upper limb, including shoulder
C49.12	Malignant neoplasm of connective and soft tissue of left upper limb, including shoulder

C49.20	Malignant neoplasm of connective and soft tissue of unspecified lower limb, including hip
C49.21	Malignant neoplasm of connective and soft tissue of right lower limb, including hip
C49.22	Malignant neoplasm of connective and soft tissue of left lower limb, including hip
C49.8	Malignant neoplasm of overlapping sites of connective and soft tissue
C49.9	Malignant neoplasm of connective and soft tissue, unspecified
C83.50	Lymphoblastic (diffuse) lymphoma, unspecified site
C83.51	Lymphoblastic (diffuse) lymphoma, lymph nodes of head, face, and neck
C83.52	Lymphoblastic (diffuse) lymphoma, intrathoracic lymph nodes
C83.53	Lymphoblastic (diffuse) lymphoma, intra-abdominal lymph nodes
C83.54	Lymphoblastic (diffuse) lymphoma, lymph nodes of axilla and upper limb
C83.55	Lymphoblastic (diffuse) lymphoma, lymph nodes of inguinal region and lower limb
C83.56	Lymphoblastic (diffuse) lymphoma, intrapelvic lymph nodes
C83.57	Lymphoblastic (diffuse) lymphoma, spleen
C83.58	Lymphoblastic (diffuse) lymphoma, lymph nodes of multiple sites
C83.59	Lymphoblastic (diffuse) lymphoma, extranodal and solid organ sites
C91.00	Acute lymphoblastic leukemia not having achieved remission
C91.01	Acute lymphoblastic leukemia, in remission
C91.02	Acute lymphoblastic leukemia, in relapse
C92.10	Chronic myeloid leukemia, BCR/ABL-positive, not having achieved remission
C92.11	Chronic myeloid leukemia, BCR/ABL-positive, in remission
C92.12	Chronic myeloid leukemia, BCR/ABL-positive, in relapse
C94.8	Other specified leukemias
C94.80	Other specified leukemias not having achieved remission
C94.81	Other specified leukemias, in remission
C94.82	Other specified leukemias, in relapse
C95.1	Chronic leukemia of unspecified cell type
C95.10	Chronic leukemia of unspecified cell type not having achieved remission
C95.11	Chronic leukemia of unspecified cell type, in remission
C95.12	Chronic leukemia of unspecified cell type, in relapse
C96.Z	Other specified malignant neoplasms of lymphoid, hematopoietic and related tissue
C96.9	Malignant neoplasm of lymphoid, hematopoietic and related tissue, unspecified
D48.1	Neoplasm of uncertain behavior of connective and other soft tissue
Z85.820	Personal history of malignant melanoma of skin
Z85.831	Personal history of malignant neoplasm of soft tissue

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC