



Skyrizi[®] (risankizumab-rzaa) (Intravenous)

Document Number: MH-0671

Last Review Date: 07/02/2024 Date of Origin: 07/05/2022 Dates Reviewed: 07/2022, 10/2022, 08/2023, 07/2024 Customization Dates: 08/08/2023, 07/02/2024 Effective Dates: 08/08/2023, 07/02/2024

I. Length of Authorization

Coverage will be provided for 9 weeks (63 days) and cannot be renewed.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

• Skyrizi carton containing one 600 mg/10 mL single-dose vial: 2 vials at Weeks 0, 4 & 8 (6 vials total)

B. Max Units (per dose and over time) [HCPCS Unit]:

- Crohn's Disease
 - Induction dose: 600 billable units (600 mg) at Week 0, 4, & 8
- Ulcerative Colitis
 - Induction dose: 1200 billable units (1200 mg) at Week 0, 4, & 8

III. Initial Approval Criteria¹

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; AND
- Physician has assessed baseline disease severity utilizing an objective measure/tool; AND
- Patient is up to date with all age-appropriate vaccinations, in accordance with current vaccination guidelines, prior to initiating therapy; **AND**
- Patient has been evaluated and screened for the presence of latent tuberculosis (TB) infection prior to initiating treatment and will receive ongoing monitoring for the presence of TB during treatment; **AND**
- Patient does not have an active infection, including clinically important localized infections; **AND**
- Patient will not receive live vaccines during therapy; **AND**



- Patient is not on concurrent treatment with another IL-inhibitor, TNF-inhibitor, biologic response modifier or other non-biologic immunomodulating agent (e.g., apremilast, tofacitinib, baricitinib, upadacitinib, abrocitinib, deucravacitinib, etc.); **AND**
- Baseline liver enzymes and bilirubin levels have been obtained prior to initiating therapy; AND

Crohn's Disease (CD) † 1,25-29

- Documented moderate to severe active disease; AND
- Documented failure, contraindication, or ineffective response at maximum tolerated doses to a minimum 3-month trial of corticosteroids or immunomodulators (e.g., azathioprine, 6-mercaptopurine, or methotrexate, etc.)

Ulcerative Colitis † 1,30-37

- Documented moderate to severe active disease; AND
 - Documented failure or ineffective response to a minimum 3-month trial of conventional therapy [aminosalicylates, corticosteroids or immunomodulators (e.g., azathioprine, 6mercaptopurine, methotrexate, etc.] at maximum tolerated doses, unless there is a contraindication or intolerance to use; OR
 - Documented failure, contraindication, or ineffective response at maximum tolerated doses to a minimum 3-month trial of a TNF modifier such as adalimumab, golimumab, or infliximab

FDA Approved Indication(s); Compendia Recommended Indication(s); Orphan Drug

IV. Renewal Criteria¹

Coverage cannot be renewed.

V. Dosage/Administration¹

Indication	Dose	
Crohn's Disease	Induction:Administer 600 mg intravenously at Week 0, Week 4, and Week 8.Maintenance:Administer 180 mg or 360 mg subcutaneously at Week 12 andevery 8 weeks thereafter (*NOTE: refer to criteria for self-administration under applicable benefit)	
Ulcerative Colitis	Induction: Administer 1200 mg intravenously at Week 0, Week 4, and Week 8. Maintenance: Administer 180 mg or 360 mg subcutaneously at Week 12 and every 8 weeks thereafter (*NOTE: refer to criteria for self-administration under applicable benefit)	

VI. Billing Code/Availability Information

HCPCS Code:

		SKYRIZI [®] IV (risankizumab-rzaa) Prior Auth Criteria
Page 2	I.	Proprietary Information. Restricted Access – Do not disseminate or copy without approval.
		©2024, Magellan Rx Management



- J2327 Injection, risankizumab-rzaa, intravenous, 1 mg; 1 billable unit = 1 mg NDC(s):
- Skyrizi carton containing one 600 mg/10 mL single-dose vial: 00074-5015-xx

VII. References

- 1. Skyrizi [package insert]. North Chicago, IL; AbbVie, Inc.; June 2024. Accessed June 2024.
- 2. Hsu S, Papp KA, Lebwohl MG, et al. Consensus guidelines for the management of plaque psoriasis. Arch Dermatol. 2012 Jan;148(1):95-102.
- 3. Menter A, Gottlieb A, Feldman SR, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 1. Overview of psoriasis and guidelines of care for the treatment of psoriasis with biologics. J Am Acad Dermatol. 2008 May;58(5):826-50.
- Gottlieb A, Korman NJ, Gordon KB, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 2. Psoriatic arthritis: overview and guidelines of care for treatment with an emphasis on the biologics. J Am Acad Dermatol 2008 May;58(5):851-64.
- National Institute for Health and Care Excellence. NICE 2008. Infliximab for the treatment of adults with psoriasis. Published 23 January 2008. Technology Appraisal Guidance [TA134]. <u>https://www.nice.org.uk/guidance/ta134/resources/infliximab-for-the-treatment-ofadults-with-psoriasis-pdf-82598193811141</u>.
- Smith CH, Jabbar-Lopez ZK, Yiu ZK, et al. British Association of Dermatologists guidelines for biologic therapy for psoriasis 2017. Br J Dermatol. 2017 Sep;177(3):628-636. Doi: 10.1111/bjd.15665.
- Armstrong AW, Siegel MP, Bagel J, et al. From the Medical Board of the National Psoriasis Foundation: Treatment targets for plaque psoriasis. J Am Acad Dermatol. 2017 Feb; 76(2):290-298. Doi: 10.1016/j.jaad.2016.10.017.
- Gordon KB, Strober B, Lebwohl M, et al. Efficacy and safety of risankizumab in moderateto-severe plaque psoriasis (UltIMMa-1 and UltIMMa-2): results from two double-blind, randomised, placebo-controlled and ustekinumab-controlled phase 3 trials. Lancet. 2018 Aug 25;392(10148):650-661. Doi: 10.1016/S0140-6736(18)31713-6. Epub 2018 Aug 7.
- Langley R, Blauvelt A, Gooderham M, et al. Efficacy and safety of continuous Q12W risankizumab versus treatment withdrawal: results from the phase 3 IMMhance trial [poster no. 10093]. In: AAD Annual Meeting. 2019.
- 10. Reich K, Gooderham M, Thaçi D, et al. Efficacy and safety of continuous risankizumab or switching from adalimumab to risankizumab treatment in patients with moderate-to-severe plaque psoriasis: results from the phase 3 IMMvent trial [poster no. 10218]. In: AAD Annual Meeting. 2019.
- 11. Richard EG. (2022). Psoralen plus ultraviolet A (PUVA) photochemotherapy. In Elmets CA, Corona R (Eds.), *UptoDate*. Lasted updated: Dec 01, 2022; Accessed on: July 20, 2023. Available from <u>https://www.uptodate.com/contents/psoralen-plus-ultraviolet-a-puva-</u>



 $\label{eq:photochemotherapy} photochemotherapy?search=Psoralen%20plus%20ultraviolet%20A%20(PUVA)%20photochemotherapy&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1.$

- 12. Hoönigsmann H. (2023). UVB therapy (broadband and narrowband). In Elmets CA, Corona R (Eds.), *UptoDate*. Last updated: Jan 18, 2023; Accessed on July 20, 2023. Available from <a href="https://www.uptodate.com/contents/uvb-therapy-broadband-and-narrowband?search=UVB%20therapy%20(broadband%20and%20narrowband&source=search-result&selectedTitle=1~80&usage_type=default&display_rank=1.
- 13. American Academy of Dermatology Work Group. Guidelines of care for the management of psoriasis and psoriatic arthritis: section 6. Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. J Am Acad Dermatol. 2011 Jul;65(1):137-74.
- 14. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. J Am Acad Dermatol 2019; 80:1029. https://doi.org/10.1016/j.jaad.2018.11.057.
- 15. Smith CH, Yiu ZZN, Bale T, et al; British Association of Dermatologists' Clinical Standards Unit. British Association of Dermatologists guidelines for biologic therapy for psoriasis 2020: a rapid update. Br J Dermatol. 2020 Oct;183(4):628-637. doi: 10.1111/bjd.19039.
- 16. National Institute for Health and Care Excellence. NICE 2017. Psoriasis: assessment and management. Published 24 October 2012. Clinical guideline [CG153]. https://www.nice.org.uk/guidance/CG153.
- 17. Menter A, Gelfand JM, Connor C, et al. Joint American Academy of Dermatology-National Psoriasis Foundation guidelines of care for the management of psoriasis with systemic nonbiologic therapies. J Am Acad Dermatol 2020; 82:1445.
- National Institute for Health and Care Excellence (NICE). Risankizumab for treating moderate to severe plaque psoriasis. Technology appraisal guidance. Published: 21 August 2019. www.nice.org.uk/guidance/ta596.
- 19. American Academy of Dermatology Work Group. Guidelines of care for the management of psoriasis and psoriatic arthritis: section 6. Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. J Am Acad Dermatol. 2011 Jul;65(1):137-74.
- 20. Gossec L, Baraliakos X, Kerschbaumer A, et al. EULAR recommendations for the management of psoriatic arthritis with pharmacological therapies: 2019 update. Ann Rheum Dis. 2020 Jun;79(6):700-712. doi: 10.1136/annrheumdis-2020-217159.
- 21. National Institute for Health and Care Excellence. NICE 2017. Certolizumab pegol and secukinumab for treating active psoriatic arthritis after inadequate response to DMARDs. Published 24 May 2017. Technology Appraisal Guidance [TA445].
- 22. Kristensen LE, Keiserman M, Papp K, et al. Efficacy and safety of risankizumab for active psoriatic arthritis: 24-week results from the randomised, double-blind, phase 3 KEEPsAKE 1 trial. Ann Rheum Dis. 2022 Feb;81(2):225-231. doi: 10.1136/annrheumdis-2021-221019.



- 23. Östör A, Van den Bosch F, Papp K, et al. Efficacy and safety of risankizumab for active psoriatic arthritis: 24-week results from the randomised, double-blind, phase 3 KEEPsAKE 2 trial. Ann Rheum Dis. 2021 Nov 23:annrheumdis-2021-221048. doi: 10.1136/annrheumdis-2021-221048.
- 24. National Institute for Health and Care Excellence. NICE 2013. Psoriasis. Published 06 August 2013. Quality standard [QS40]. https://www.nice.org.uk/guidance/qs40.
- 25. D'Haens G, Panaccione R, Baert F, et al. Risankizumab as induction therapy for Crohn's disease: results from the phase 3 ADVANCE and MOTIVATE induction trials. Lancet. 2022 May 28;399(10340):2015-2030. doi: 10.1016/S0140-6736(22)00467-6.
- 26. Lichtenstein GR, Loftus EV, Isaacs KI, et al. ACG Clinical Guideline: Management of Crohn's Disease in Adults. Am J Gastroenterol 2018; 113:481–517; doi: 10.1038/ajg.2018.27.
- 27. Torres J, Bonovas S, Doherty G, et al. European Crohn's and Colitis Organisation [ECCO] Guidelines on Therapeutics in Crohn's Disease: Medical Treatment. Journal of Crohn's and Colitis, 2020, 4–22 doi:10.1093/ecco-jcc/jjz180.
- 28. Terdiman JP, Gruss CB, Heidelbaugh JJ, et al. American Gastroenterological Association Institute Guideline on the use of Thiopurines, Methotrexate, and Anti-TNF-α Biologic Drugs for the Induction and Maintenance of Remission in Inflammatory Crohn's Disease. Gastroenterology 2013;145:1459-1463.
- 29. National Institute for Health and Care Excellence. NICE 2019. Crohn's Disease: Management. Published 03 May 2019. NICE Guideline [NG129]. https://www.nice.org.uk/guidance/ng129.
- 30. Harbord M, Eliakim R, Bettenworth D, et al. Third European Evidence-based Consensus on Diagnosis and Management of Ulcerative Colitis. Part 2: Current Management. J Crohns Colitis. 2017 Jan 28. doi: 10.1093/ecco-jcc/jjx009.
- Lewis JD, Chuai S, Nessel L, et al. Use of the Non-invasive Components of the Mayo Score to Assess Clinical Response in Ulcerative Colitis. Inflamm Bowel Dis. 2008 Dec; 14(12): 1660–1666. doi: 10.1002/ibd.20520.
- 32. Paine ER. Colonoscopic evaluation in ulcerative colitis. Gastroenterol Rep (Oxf). 2014 Aug; 2(3): 161–168.
- 33. Walsh AJ, Bryant RV, Travis SPL. Current best practice for disease activity assessment in IBD. Nature Reviews Gastroenterology & Hepatology 13, 567–579 (2016) doi:10.1038/nrgastro.2016.128.
- 34. Feuerstein JD, Isaacs KL, Schneider Y, et al. AGA Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis. Gastroenterology. 2020;158(5):1450-1461. doi:10.1053/j.gastro.2020.01.006.
- 35. Raine T, Bonovas S, Burisch J, et al. ECCO Guidelines on therapeutics in ulcerative colitis: medical treatment. J Crohns Colitis. 2022 Jan 28. 16 (1):2-17. Doi: 10.1093/ecco-jcc/jjab178.
- 36. Louis E, Panaccione R, Parkes G, et al. S4 Risankizumab Induction Therapy in Patients With Moderately to Severely Active Ulcerative Colitis: Efficacy and Safety in the



Randomized Phase 3 INSPIRE Study. The American Journal of Gastroenterology 118(12S):p S2, December 2023. | DOI: 10.14309/01.ajg.0000995752.19097.c8

37. Louis E, Panaccione R, Parkes G, et al. OP06 Risankizumab Maintenance Therapy in Patients With Moderately to Severely Active Ulcerative Colitis: Efficacy and Safety in the Randomised Phase 3 COMMAND Study, Journal of Crohn's and Colitis, Volume 18, Issue Supplement_1, January 2024, Pages i10–i12, https://doi.org/10.1093/ecco-jcc/jjad212.0006

ICD-10	ICD-10 Description	
Codes		
K50.00	Crohn's disease of small intestine without complications	
K50.011	Crohn's disease of small intestine with rectal bleeding	
K50.012	Crohn's disease of small intestine with intestinal obstruction	
K50.013	Crohn's disease of small intestine with fistula	
K50.014	Crohn's disease of small intestine with abscess	
K50.018	Crohn's disease of small intestine with other complication	
K50.019	Crohn's disease of small intestine with unspecified complications	
K50.10	Crohn's disease of large intestine without complications	
K50.111	Crohn's disease of large intestine with rectal bleeding	
K50.112	Crohn's disease of large intestine with intestinal obstruction	
K50.113	Crohn's disease of large intestine with fistula	
K50.114	Crohn's disease of large intestine with abscess	
K50.118	Crohn's disease of large intestine with other complication	
K50.119	Crohn's disease of large intestine with unspecified complications	
K50.80	Crohn's disease of both small and large intestine without complications	
K50.811	Crohn's disease of both small and large intestine with rectal bleeding	
K50.812	Crohn's disease of both small and large intestine with intestinal obstruction	
K50.813	Crohn's disease of both small and large intestine with fistula	
K50.814	Crohn's disease of both small and large intestine with abscess	
K50.818	Crohn's disease of both small and large intestine with other complication	
K50.819	Crohn's disease of both small and large intestine with unspecified complications	
K50.90	Crohn's disease, unspecified, without complications	
K50.911	Crohn's disease, unspecified, with rectal bleeding	
K50.912	Crohn's disease, unspecified, with intestinal obstruction	
K50.913	Crohn's disease, unspecified, with fistula	
K50.914	Crohn's disease, unspecified, with abscess	

Appendix 1 – Covered Diagnosis Codes

SKYRIZI® IV (risankizumab-rzaa) Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval. ©2024, Magellan Rx Management

ICD-10	ICD-10 Description	
Codes		
K50.918	Crohn's disease, unspecified, with other complication	
K50.919	Crohn's disease, unspecified, with unspecified complications	
K51.00	Ulcerative (chronic) pancolitis without complications	
K51.011	Ulcerative (chronic) pancolitis with rectal bleeding	
K51.012	Ulcerative (chronic) pancolitis with intestinal obstruction	
K51.013	Ulcerative (chronic) pancolitis with fistula	
K51.014	Ulcerative (chronic) pancolitis with abscess	
K51.018	Ulcerative (chronic) pancolitis with other complication	
K51.019	Ulcerative (chronic) pancolitis with unspecified complications	
K51.20	Ulcerative (chronic) proctitis without complications	
K51.211	Ulcerative (chronic) proctitis with rectal bleeding	
K51.212	Ulcerative (chronic) proctitis with intestinal obstruction	
K51.213	Ulcerative (chronic) proctitis with fistula	
K51.214	Ulcerative (chronic) proctitis with abscess	
K51.218	Ulcerative (chronic) proctitis with other complication	
K51.219	Ulcerative (chronic) proctitis with unspecified complications	
K51.30	Ulcerative (chronic) rectosigmoiditis without complications	
K51.311	Ulcerative (chronic) rectosigmoiditis with rectal bleeding	
K51.312	Ulcerative (chronic) rectosigmoiditis with intestinal obstruction	
K51.313	Ulcerative (chronic) rectosigmoiditis with fistula	
K51.314	Ulcerative (chronic) rectosigmoiditis with abscess	
K51.318	Ulcerative (chronic) rectosigmoiditis with other complication	
K51.319	Ulcerative (chronic) rectosigmoiditis with unspecified complications	
K51.50	Left sided colitis without complications	
K51.511	Left sided colitis with rectal bleeding	
K51.512	Left sided colitis with intestinal obstruction	
K51.513	Left sided colitis with fistula	
K51.514	Left sided colitis with abscess	
K51.518	Left sided colitis with other complication	
K51.519	Left sided colitis with unspecified complications	
K51.80	Other ulcerative colitis without complications	
K51.811	Other ulcerative colitis with rectal bleeding	
K51.812	Other ulcerative colitis with intestinal obstruction	

SKYRIZI® IV (risankizumab-rzaa) Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval. ©2024, Magellan Rx Management

ICD-10	ICD-10 Description	
Codes		
K51.813	Other ulcerative colitis with fistula	
K51.814	Other ulcerative colitis with abscess	
K51.818	Other ulcerative colitis with other complication	
K51.819	Other ulcerative colitis with unspecified complications	
K51.90	Ulcerative colitis, unspecified, without complications	
K51.911	Ulcerative colitis, unspecified with rectal bleeding	
K51.912	Ulcerative colitis, unspecified with intestinal obstruction	
K51.913	Ulcerative colitis, unspecified with fistula	
K51.914	Ulcerative colitis, unspecified with abscess	
K51.918	Ulcerative colitis, unspecified with other complication	
K51.919	Ulcerative colitis, unspecified with unspecified complications	
R19.7	Diarrhea, unspecified	

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: https://www.cms.gov/medicare-coverage-database/search.aspx. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions				
Jurisdiction	Applicable State/US Territory	Contractor		
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC		
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC		
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)		
6	MN, WI, IL	National Government Services, Inc. (NGS)		
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.		
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)		
N (9)	FL, PR, VI	First Coast Service Options, Inc.		
J (10)	TN, GA, AL	Palmetto GBA		
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA		

SKYRIZI® IV (risankizumab-rzaa) Prior Auth Criteria

Medicare Part B Administrative Contractor (MAC) Jurisdictions				
Jurisdiction	Applicable State/US Territory	Contractor		
	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.		
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)		
15	KY, OH	CGS Administrators, LLC		

Page 9

