

Talvey™ (talquetamab-tgvs) (Subcutaneous)

Document Number: IC-0722

Last Review Date: 09/05/2023 Date of Origin: 09/05/2023 Dates Reviewed: 09/2023

I. Length of Authorization

After the initial hospital administration of three step-up titration doses, coverage will be provided for six months and may be renewed.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- Talvey 3 mg/1.5 mL solution for injection in a single-dose vial: 3 vials per week
- Talvey 40 mg/1 mL solution for injection in a single-dose vial: 3 vial per week

B. Max Units (per dose and over time) [HCPCS Unit]:

- Titration: 3 mg on day one, 9 mg on day four and 40 mg on day seven
- Maintenance: 40 mg weekly thereafter

III. Initial Approval Criteria ¹

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; AND
- Prescribers are enrolled in the TECVAYLI-TALVEY REMS® program; AND
- Used as continuation therapy following inpatient administration of all the step-up doses;
 AND
- Patient had an absence of unacceptable toxicity while on inpatient administration of stepup titration doses; AND

Universal Criteria 1

- Patient does not have an active infection, including clinically important localized infections;
 AND
- Prophylaxis for infection will be followed according to local guidelines; AND
- Patient does not have active CNS involvement or clinical signs of meningeal involvement of multiple myeloma; **AND**



- Patient has not had an allogenic stem cell transplant within the previous six months or an autologous stem cell transplant within the previous 12 weeks; **AND**
- Patient weight and signs of oral and skin toxicity will be monitored at baseline and periodically during therapy; **AND**

Multiple Myeloma † ‡ Φ 1-5

- Used as single-agent therapy; AND
- Patient has relapsed or refractory disease; AND
- Patient has received at least four (4) prior therapies, including a proteasome inhibitor (e.g., bortezomib, etc.), an immunomodulatory agent (e.g., lenalidomide, thalidomide, etc.) and an anti-CD38 monoclonal antibody (e.g., daratumumab, isatuximab, etc.)

† FDA Approved Indication(s); ‡ Compendia Approved Indication(s); **Φ** Orphan Drug

IV. Renewal Criteria ¹

Coverage may be renewed based upon the following criteria:

- Patient continues to meet the indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; AND
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: severe oral toxicity and weight loss, serious life-threatening infections, severe cytopenias, severe dermatologic toxicity, hepatotoxicity, neurologic toxicity including Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS), cytokine release syndrome (CRS), etc.

V. Dosage/Administration ¹

Indication	Dos	Dose				
Multiple Myeloma	The recommended dosage is administered subcutaneously by a healthcare provider, once or bi-weekly (every 2 weeks) until disease progression or unacceptable toxicity.					
		Dosing schedule	Day	Dose		
		Step-up dosing schedule ^a	Day 1	Step-up dose 1	0.01 mg/kg	
			Day 4 ^b	Step-up dose 2	0.06 mg/kg	
			Day 7 ^c	First treatment dose	0.4 mg/kg	
		Weekly dosing schedule ^a	One week after first treatment dose and weekly thereafter	Subsequent treatment doses	0.4 mg/kg once weekly	



- b Dose may be administered between 2 to 4 days after the previous dose and may be given up to 7 days after the previous dose to allow for
- c Maintain a minimum of 6 days between weekly doses

Dosing schedule	Day	Dose	
	Day 1	Step-up dose 1	0.01 mg/kg
Step-up dosing	Day 4 ^b	Step-up dose 2	0.06 mg/kg
schedule ^a	Day 7 ^b	Step-up dose 3	0.4 mg/kg
	Day 10 ^c	First treatment dose	0.8 mg/kg
Biweekly (every 2 weeks) dosing schedule ^a	Two weeks after first treatment dose and 2 weeks thereafter ^d	Subsequent treatment doses	0.8 mg/kg every 2 weeks

- b Dose may be administered between 2 to 4 days after the previous dose and may be given up to 7 days after the previous dose to allow for
- c Dose may be administered between 2 to 7 days after step-up dose 3.
- d Maintain a minimum of 12 days between biweekly (every 2 weeks) doses.

Note: Administer Talvey subcutaneously according to the step up dosing schedule noted above to reduce the incidence and severity of cytokine release syndrome (CRS). Due to the risk of CRS and neurologic toxicity, including ICANS, patients should be hospitalized for 48 hours after administration of all doses within the Talvey step-up dosing schedule

VI. **Billing Code/Availability Information**

HCPCS Code:

J9999 – Not otherwise classified, antineoplastic drugs

NDC:

- Talvey 3 mg/1.5 mL solution for injection in a single-dose vial: 57894-0469-xx
- Talvey 40 mg/1 mL solution for injection in a single-dose vial: 57894-0470-xx

VII. References

- 1. Talvey [package insert]. Horsham, PA; Janssen Biotech, Inc.; August 2023. Accessed August 2023.
- 2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for talquetamab. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed August 2023.



- 3. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Multiple Myeloma Version 3.2023. National Comprehensive Cancer Network, 2023. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc." To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed August 2023.
- 4. BGM Durie, J-L Harousseau, J S Miguel, et al on behalf of the International Myeloma Working Group. International uniform response criteria for multiple myeloma. Leukemia. Sep; 20(9):1467-73.
- 5. Schinke CD, Touzeau C, Minnema MC, et al. Pivotal phase 2 MonumenTAL-1 results of talquetamab (tal), a GPRC5DxCD3 bispecific antibody (BsAb), for relapsed/refractory multiple myeloma (RRMM). Journal of Clinical Oncology 2023 41:16_suppl, 8036-8036.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description	
C90.00	Multiple myeloma not having achieved remission	
C90.02	Multiple myeloma in relapse	
C90.10	Plasma cell leukemia not having achieved remission	
C90.12	Plasma cell leukemia in relapse	
C90.20	Extramedullary plasmacytoma not having achieved remission	
C90.22	Extramedullary plasmacytoma in relapse	
C90.30	Solitary plasmacytoma not having achieved remission	
C90.32	Solitary plasmacytoma in relapse	
Z85.79	Personal history of other malignant neoplasms of lymphoid, hematopoietic and related tissues	

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: https://www.cms.gov/medicare-coverage-database/search.aspx. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions					
Jurisdiction	Applicable State/US Territory	Contractor			
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC			
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC			
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)			



Medicare Part B Administrative Contractor (MAC) Jurisdictions				
Jurisdiction	Applicable State/US Territory	Contractor		
6	MN, WI, IL	National Government Services, Inc. (NGS)		
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.		
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)		
N (9)	FL, PR, VI	First Coast Service Options, Inc.		
J (10)	TN, GA, AL	Palmetto GBA, LLC		
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC		
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.		
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)		
15	KY, OH	CGS Administrators, LLC		

